

# Non-operative Clavicle Fracture

## PHASE I: Protect clavicle (*Injury to 2 weeks*)

- Sling at all times
- No lifting with injured arm
- Motion: No shoulder ROM
- Elbow and forearm exercises, ball squeeze exercise
- Strengthening: No resistive exercises/activities

## PHASE II: Advance ROM (*2-6 weeks*)

- Sling: for comfort
- Motion: Begin gentle PROM exercises followed by AAROM and AROM in all planes to pain tolerance
- Strengthening: Begin gentle Theraband resistive exercises at 4 weeks

## PHASE III: Restore function (*6-12 weeks*)

- Sling: Discontinue sling at all times
- Motion: goal for full motion by week 12
- Strengthening: Progress to higher weights and sports specific training at week 10
- Return to sports 3-6 months from surgery

**Gunnison** | 112 W. Spencer Ave., Suite A | Gunnison, CO 81230 | Phone: 970-641-6788 | Fax: 970-641-0282

**Crested Butte** | 405 Elk Ave. | Crested Butte, CO 81224 | Phone: 970-641-6788 | Fax: 970-641-0282

**Telluride** | 500 W. Pacific Ave. | Telluride, CO 81435 | Phone: 970-641-6788 | Fax: 970-641-0282