

**Adam Lindsay, MD, M.S.**

VSON Alpine - Gunnison, Crested Butte, Telluride  
Orthopaedic Surgery, Sports Medicine  
P) 970-641-6788 F) 866-725-4659



## General Post-Operative Instructions

### Distal Biceps Tendon Repair

#### Icing

Icing is very important for the first 7-14 days after surgery. You will be given the option to rent/purchase a cooling machine – this will be arranged before surgery. For the first 3-4 days, a bag of ice or the cooling machine should be used as frequently as possible. After 4 days, ice or cooling machine is applied for 20-minute periods, at least 4 times per day. Care must always be taken to avoid frostbite – there should be a layer of clothing or a cloth/towel between the ice and skin.

#### Nerve Block

Anesthesia will typically provide a nerve block after your surgery while you are still in the recovery room. It is your choice as to if you would like to receive the nerve block, but it is generally recommended to help with pain control. This decision is made the day of surgery after the anesthesia team explains the risks and benefits. The nerve block is an injection in the area of your neck that numbs the nerves in your arm to reduce pain after the surgery. There are 2 types of nerve blocks: single shot and continuous (nerve catheter). The single shot block typically lasts 14 to 24 hours but this is variable. The continuous nerve block involves the placement of a catheter to allow medication to numb your arm for several days. It is not uncommon to lose the ability to move your arm and hand after the block (often described as a dead arm) – this will resolve as the block wears off.

#### Sling Use

A sling will be fitted and placed in the operating room for your comfort. You may release the strap on the sling to straighten and bend the elbow 3 to 4 times per day starting the day after surgery. During the first 2 weeks, you may remove your sling at home during the day but need to wear the sling in public (around other people) and when sleeping. After 2 weeks, the sling can be removed at all times.

**Adam Lindsay, MD, M.S.**  
VSON Alpine - Gunnison, Crested Butte, Telluride  
Orthopaedic Surgery, Sports Medicine  
P) 970-641-6788 F) 866-725-4659



## **Physical Therapy**

Physical therapy is to be scheduled to start 2 weeks after surgery. It is important to select your physical therapy location and call ahead of time to setup an appointment as many

**Adam Lindsay, MD, M.S.**

VSON Alpine - Gunnison, Crested Butte, Telluride  
Orthopaedic Surgery, Sports Medicine  
P) 970-641-6788 F) 866-725-4659



physical therapy locations are scheduled out for 1-2 weeks.

You may go to any physical therapy location you wish, but it is ideal to pick one that is close to your home or work. You will be provided a prescription for physical therapy and a rehabilitation protocol at your pre-operative appointment. Please bring these forms to your therapist. If you forget or lose them, you can find them at [www.lindsayssportsmed.com](http://www.lindsayssportsmed.com) under the 'Therapy Protocols' section. Again, you are encouraged to call before or soon after surgery and schedule an appointment to start 2 weeks after surgery.

## Incision Care

After surgery, you will have a splint on your arm. This will stay in place until I remove it at your first post-operative appointment (10-14 days). Sutures will also be removed at your first post-operative appointment. You will be provided a hinged elbow brace at your first post-operative appointment that is to be worn at all times. You may shower after surgery with the splint in place.

The splint **CANNOT** get wet during your shower. Wrap the splint with a dry towel and place your arm in a large trash bag. **DO NOT** get into a bathtub, pool, or spa until your sutures are removed and your incisions are completely healed to lower the chance of skin infection. Always wash your hands before touching your incisions. **DO NOT** use any creams or ointments on your incisions.

**Adam Lindsay, MD, M.S.**

VSON Alpine – Gunnison, Crested Butte, Telluride  
Orthopaedic Surgery, Sports Medicine  
P) 970-641-6788 F) 866-725-4659



Please contact my office immediately if you notice any of the following as these could be a sign of an infection:

1. Excessive fluid leaking from the incisions after 3 days
2. A foul odor from the incisions
3. Any redness or warmth around the incisions
4. Significant increase in pain.

In addition, please check your temperature if you begin to feel ill, warm, or if you have chills. Contact my office immediately if your temperature is above 101.5 degrees or you think you may have an infection anywhere in your body. It is common to have a low-grade temperature within the first week of surgery. You should drink fluids and breathe deeply.

## **Swelling and Bruising**

It is common to have swelling and/or bruising after surgery. This is caused by bleeding and is expected. The bruising may start out black-red-purple and will change to a yellowish-green color as it fades over a few weeks. The bruising may go down the arm. You may also have some swelling in your hand which should go away; squeezing a ball or making a fist repeatedly will help with this. In time, the swelling will go away.

## **Discomfort**

You may need help with your daily activities, so it is a good idea to have family and friends around to help you. You will have some mild to moderate elbow discomfort. You will be provided a prescription for pain pills after the surgery. Please take your pills as directed and remember to use ice or your cooling device to help reduce pain. If you have pain that your pain pills do not take care of, please contact my office.

## **Nausea and Vomiting**

It is common to have nausea and/or vomiting for the first 24 hours after surgery. You have been provided with a medication (Ondansetron) that will help with nausea. Please contact my office about any nausea/vomiting that does not go away. You may eat whatever you wish, however, it is suggested that you start with fluids and bland food before moving to a regular diet.

**Adam Lindsay, MD, M.S.**

VSON Alpine - Gunnison, Crested Butte, Telluride  
Orthopaedic Surgery, Sports Medicine  
P) 970-641-6788 F) 866-725-4659



## **Your Medications**

You should start taking all your regular medications right after surgery. If there are exceptions to this, you will be informed. If you have questions contact my office.

## **Blood Clots**

Surgery may slow the blood flow in your legs, which might (rarely) cause a blood clot to form in the leg. If a clot were to form, the leg is usually painful and swollen. Blood clots can be serious and, if you have one, you will need to go to the hospital. Walking regularly early after surgery can prevent blood clots. Moving the ankle frequently and rising up on your toes can lower the risk of this problem. If you are not already on a blood thinner, I would like you to take a full-strength Aspirin (325mg) once daily for 3 weeks - this can help prevent a blood clot. If you were taking a blood thinner before surgery, I will let you know when you can start taking this again after surgery (most often the day after surgery).

### **Please contact your doctor right away if:**

You have swelling, tenderness, pain, warmth or redness in your thigh, calf, or ankle.

### **Call 911 if you have signs that might mean a blood clot that has moved to your lungs:**

- Chest Pain
- Trouble Breathing
- Rapid Breathing
- Sweating
- Confusion